



# Determining puma habitat suitability in the Eastern USA

Veronica Yovovich<sup>1,2</sup> · Nathaniel Robinson<sup>1,3</sup> · Hugh Robinson<sup>1,4</sup> · Michael J. Manfredo<sup>5</sup> · Shelby Perry<sup>6</sup> · Jeremy T. Bruskotter<sup>7</sup> · John A. Vucetich<sup>8</sup> · Luis Aníbal Solórzano<sup>9</sup> · Lydia A. Roe<sup>6</sup> · Alison Lesure<sup>10</sup> · Jamie Robertson<sup>1</sup> · Tom Butler<sup>6</sup> · L. Mark Elbroch<sup>1</sup>

Received: 16 June 2022 / Revised: 14 November 2022 / Accepted: 29 November 2022 /  
Published online: 3 January 2023  
© The Author(s) 2022

## Abstract

Pumas (*Puma concolor*) were eliminated from most of the eastern USA a century ago. In the past couple of decades, their recovery in the West has increased puma dispersal into the Midwest, with some individuals even traveling to the East Coast. We combined published expert opinion data and a habitat suitability index in an analysis that identified 17 areas in the Upper Midwest, Ozarks, Appalachia, and New England that could potentially host puma populations in the future. Thirteen of these were larger than 10,000 km<sup>2</sup> and so likely to ensure a puma population's long-term genetic health. Further, we quantified patch size, human density, livestock density, percent public land, and a sociocultural index reflecting wildlife values for comparing patches, as well as present a summary of current legislation relevant to puma management in the East. Our work may be useful in identifying suitable areas to restore pumas based not only on the quality of their biophysical habitat, but also on social values conducive to puma-human coexistence.

**Keywords** Habitat connectivity · Habitat suitability · Landscape connectivity · *Puma concolor* · Range expansion · Recolonize

## Introduction

One hundred fifty years ago, pumas (*Puma concolor*), also called cougars, mountain lions, and Florida panthers, roamed nearly every habitat in the Americas from the East to West coast and southern Canada to Chile (Culver et al. 2000; McCollough 2011). However, when European settlers spread across North America, they viewed carnivores as competitors for game, livestock and other resources (Gill 2010). As a result, they conducted wide-scale carnivore eradication efforts (Danz 1999). Early settlers also practiced unrestricted resource extraction, resulting in deforestation and precipitous declines

---

Communicated by Akihiro Nakamura.

✉ Veronica Yovovich  
yvovovich@panthera.org

Extended author information available on the last page of the article

of ungulate prey that supported large carnivores (Gill 2010). With the establishment of an independent USA, these practices evolved into federal and state bounty programs by the late 19th century, incentivizing the removal of pumas and other carnivores (Feldman 2007; Gill 2010). By the early 1900s, pumas were functionally extinct in the eastern two-thirds of the country, except southern Florida (Culver et al. 2000; McCollough 2011).

In the early 20th century, the USA transformed its wildlife management to restrict hunting and protect game species humans valued, which serendipitously protected the prey species needed to recover large carnivore populations as well (Gill 2010). In the 1970s, people increasingly perceived carnivores as important components of ecosystems, and pumas were given some level of protection across most of the West (Mattson and Clark 2010; Pritchard 2021). Since that time, western puma populations have recovered remarkably. Though pumas in the east are the same subspecies as those found in western North America (Culver et al. 2000), these protections did not have the same effect on eastern populations. The largest block of historic puma range in all the Americas that does not currently support breeding puma populations comprises the Midwest and Eastern USA (Nielsen et al. 2017; Fig. 1).

Since pumas have been extirpated from much of their previous distribution, they have been unable to perform important functions that contribute to ecological resilience (Barry et al. 2019; LaBarge et al. 2022). In addition, the absence of pumas has consequences for human health and wellbeing as well. For example, pumas mitigate human-deer collisions on roadways, saving human lives and millions of dollars in costs (Gilbert et al. 2021). There is also evidence that suggests pumas help control chronic wasting disease (Baune et al. 2021), which poses great concern for wildlife agencies and hunting communities. Reestablishing pumas in currently unoccupied habitat could help restore these important ecological relationships.

Early stages of puma recolonization have been documented in the Midwest and central-southern Canada (LaRue et al. 2012; LaRue and Nielson 2011; Glick 2014; O'Neil et al. 2014; Smith et al. 2016; Gantchoff et al. 2021; LaRue et al. 2022), spurring considerable research aimed at predicting how pumas will navigate current landscapes and where they may occur in the future (LaRue and Nielsen 2008). Pumas exhibit male-biased dispersal patterns in which males disperse greater distances than females (Sweanor et al. 2000), and therefore we expect males to appear in the East well before females (LaRue et al. 2012). For example, one young male puma dispersed from the Black Hills in South Dakota approximately 1800 miles to Connecticut, where he was killed on a highway in 2011 (Hawley et al. 2016). Several reports and papers have also explored the potential for reintroducing pumas in the East, as a conservation restoration strategy to expedite recolonization of former range (Brocke 1981; Laundre 2013).

Given the density of people and human infrastructure in the East relative to the West, an important next step in managing recolonizing carnivores is determining whether they can maintain viable populations on their own. Identifying potential habitat blocks that may host pumas in the future may also be useful in targeting proactive conflict mitigation strategies and other educational campaigns that support human coexistence with large carnivores.

Here, we combine expert opinion published in peer-reviewed literature (LaRue and Nielsen 2008) and a habitat suitability index to identify habitat patches where pumas could potentially reestablish locally self-sustaining populations in the East, without relying on dispersal and genetic rescue from nearby populations (i.e., metapopulation dynamics; Sweanor et al. 2000). The identified areas contain sufficient resources for supporting a long term puma population once established, independent of neighboring populations.



**Fig. 1** Current (shown in dark green) and historic (shown in dark green and light green combined) puma range. The largest unoccupied area, or the area with the largest opportunity for recolonization, is the eastern two-thirds of the USA. Figure adapted from IUCN Red List of Threatened Species: *Puma concolor*. (Color figure online)

## Materials and methods

### Study area

Our area of focus was the USA states bordering or east of the Mississippi River, and included Alabama, Arkansas, Connecticut, Delaware, District of Columbia, Florida, Georgia, Illinois, Indiana, Iowa, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, New Hampshire, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Rhode Island, South Carolina, Tennessee, Vermont, Virginia, West Virginia, and Wisconsin. This large area was primarily composed of two ecoregions (EPA 2022), Eastern temperate forests and Northern forests, with a small portion of tropical wet forests in southern Florida. The climate, land use, habitats, and other landscape features were highly variable across this region. Regional climate ranged from continental, with cold or frigid winters and hot, humid summers; to humid subtropical, with mild winters and warm to hot and humid summers. Habitat types were similarly variable, from boreal forest and alpine regions in New England to tallgrass prairie and temperate deciduous forest in the Midwest to pond cypress swamp and pine savanna in the South.

### Identifying puma habitat patches

Pumas have been widely studied in the USA, which has provided a wealth of information on puma habitat requirements, including their selection preferences for vegetation types, human density, road proximity, livestock use, and snow depth (Table 1). We used expert opinion found in published scientific literature to identify several landscape dynamics that influence suitable puma habitat availability, puma habitat selection, as well as puma population viability. For example, pumas are ambush predators that require vegetative structure for cover, and therefore they select against open habitats (Gray et al. 2016). Pumas also select against areas of deep snow, likely because it hinders movement and reduces ungulate prey abundance (CDFW 2015; Laundré and Hernández 2003; Poole and Mowat 2005). Though not a wilderness obligate species, pumas also select against urban and suburban areas (Burdett et al. 2010; Wilmers et al. 2013; Yovovich et al. 2020). Pumas utilize fire roads, dirt roads, and other low-speed, low-use roadways, but their movement is limited by high-speed roadways, highways, and interstates (Dickson et al. 2005; Knopff et al. 2014; Wilmers et al. 2013). Conflict as a product of carnivore-livestock interactions is among the greatest threats to carnivores worldwide, and pumas are no exception (Ripple et al. 2014). Habitat with limited livestock may be more suitable puma habitat, because it reduces the risk of puma mortality via legal depredation permits or retaliatory killing (Guerisoli et al. 2021). We selected seven ecological and geographical variables that reflect puma habitat requirements and implemented a threshold approach to delineate areas of viable habitat for each variable (Table 1). We excluded all areas that did not meet our threshold requirements for all seven variables, and then combined connected habitat into patches for further analyses (for variable preprocessing and associated metadata, see supplemental material).

Based on research on puma genetic diversity, we applied two size thresholds to our resulting patches to identify best habitat for potential puma populations in the East (Dellinger et al. 2020). First, we excluded all areas less than 6,000 km<sup>2</sup>, which some evidence suggests is the minimum area required to maintain sufficient puma genetic diversity to sustain long-term populations, and then a stricter 10,000 km<sup>2</sup> threshold, as this has been

**Table 1** Variable thresholds for puma habitat suitability to be applied to the East Coast, as defined by previous literature, and data sources used in the modeling analyses for predicting suitable patches for puma survival and persistence

Ecological assumption	Habitat variable	Threshold description	Data source	Spatial resolution	Dates
Pumas select against open habitats in North America (LaRue and Nielsen 2008; Burdett et al. 2010; Gray et al. 2016)	Land cover	Exclude agriculture, high elevation rock and grasslands	Sulla-Menashe and Friedl (2018)	500 m	2019
Pumas require structured habitat for hunting and survival (Gray et al. 2016)	Forest cover	Pumas excluded from cells with < 53% forest cover	Hansen et al. (2013)	30 m	2000–2020
Pumas follow prey, and deer and elk avoid deep snow (Laundré and Hernández, 2003; Poole and Mowat, 2005)	Maximum snow depth	Pumas (and puma prey) excluded when average winter snow depth ≥ 50 cm	SNODAS	1 km	2010–2020
Pumas excluded from areas with high housing density (Burdett et al. 2010)	Housing density	Pumas excluded when housing density ≥ 0.68 units/ha	ICLUS	90 m	2010
Pumas excluded from habitat adjacent to people (Wilmers et al. 2013; Yovovich et al. 2021)	Human development proximity	Pumas excluded from habitat ≤ 600 m from human structures	Yang et al. (2018)	30 m	2016
People excluded from habitat immediately adjacent to large highways (Knopff et al. 2014)	Highway proximity	Pumas excluded from habitat within 170 m of interstate highways and major arterials	TIGER US Census Roads	NA	2016
Pumas and livestock may have conflict that leads to lower puma survival; puma survival is higher in habitat with low livestock density (Guerisoli et al. 2021)	Livestock density	Pumas excluded from habitat with ≥ 14.5 animals/km <sup>2</sup>	Robinson et al. (2014)	10 km	2010

suggested by researchers as the size needed to ensure long-term genetic health (Dellinger et al. 2020). All spatial analyses were conducted using Google Earth Engine (Gorelick et al. 2017).

### Descriptive metrics for habitat patches


For all the patches that met either size threshold (i.e.,  $> 6,000\text{km}^2$  or  $> 10,000\text{km}^2$ ), we created a suite of metrics across which we might compare them by (1) total patch area, (2) mean sociocultural index values based on county-level projections in Manfredi et al. (2021), (3) percentage of the patch that is public land (Conservation Biology Institute 2012), (4) mean human population density (GPWw4; CIESIN 2018), and (5) mean livestock density (Robinson et al. 2014). The sociocultural index values (Manfredi et al. 2021) range from 0 to 1, and reflect a breadth of perspectives on wildlife and natural resources, from domination, where wildlife are viewed primarily as resources for human consumption and benefit, to mutualism, where wildlife are viewed as part of one's social community, having rights like humans, and deserving of care and compassion. When compared with mutualists, those with a domination orientation are less tolerant of predators and more supportive of lethal control in dealing with them. The index value represents the proportion of mutualists in a given area, or the number of people exhibiting mutualist scores divided by the total number of people. Index values closer to 0 indicate a population that leans towards a domination perspective, and values closer to 1 indicate a population that leans towards mutualism. We also reviewed the wildlife action plans, regulations, and legislation as they might apply to puma conservation for the states identified as having suitable habitat.

## Results

We identified 17 areas in the Upper Midwest, Ozarks, Appalachia, and New England that met all of the minimal habitat suitability requirements, of which 13 met our stricter 10,000  $\text{km}^2$  size requirements as well (Fig. 2; Table 2). The 17 patches identified ranged in size from 6024 to 59,462  $\text{km}^2$ , mean sociocultural index values from 0.31 to 0.548, percentage public land from 3 to 80%, mean human population density from 1.76 to 35.78 people/ $\text{km}^2$ , and mean livestock density from 1.48 to 8.83 animals/ $\text{km}^2$  (Table 2).

Light and dark green show individual patches that are adjacent, but separated by unsuitable habitat, highways in most cases. These areas are likely to have some gene flow between the two, but contain sufficiently unstable habitat to be modeled as distinct areas. Note the red circle encompassing northern Vermont and New Hampshire in Fig. 2a and the close up in Fig. 2b. This is an example of our conservative methods that likely underestimate habitat where pumas could survive. Highways divided the area in the red circle into habitat patches too small to meet our threshold sizes, however traffic volumes in northern New England are likely so low as to allow connectivity across these areas. Even if pumas do not thrive in this area, but are able to move through it, the surrounding patches are likely to benefit from increased gene flow.

Most states that contained potential puma habitat patches list pumas as extirpated, and some retain laws and regulations that will impact future puma dispersal, establishment, and management (Table 3). For example, pumas are listed as endangered under the state Endangered Species Acts of New York, Vermont, South Carolina, Georgia, and Michigan; and they are protected as a species of special concern under the state Endangered Species



**Fig. 2** **a** Overview of Potential habitat patches that could support pumas identified by our analyses (shown in light and dark green) in the eastern United States, in relation to urban areas (shown in orange). **b** Close up of identified suitable areas for puma recolonization in the New England. (Color figure online)






Fig. 2 (continued)



**Table 2** Potential puma habitat patches and habitat suitability values, listed in order of increasing size

Patch number	Area name	Patch size (km <sup>2</sup> )	Percent public land	Human density (people/km <sup>2</sup> )	Sociocultural index	Livestock density (animals/km <sup>2</sup> )
1	Ouachita Mountains	6024	80	1.76	0.31	8.83
2	Green Mountains East	6517	14	35.78	0.522	3.28
3	Michigan East	7773	45	4.76	0.346	2.93
4	Michigan West	9639	46	9.75	0.383	4.01
5	Green Mountains West	11.874	22	13.69	0.548	5.55
6	Allegheny Plateau	12.040	3	11.55	0.35	7.34
7	Catskill Mountains	12.451	15	18.34	0.427	4.77
8	Maine East	12.831	5	8.87	0.445	1.48
9	Ozark Plateau	14.341	34	8.11	0.317	8.05
10	Great Smoky South	17.099	53	19.23	0.394	4.23
11	Appalachian North	21.204	29	9.1	0.353	6.39
12	Allegheny North	21.582	42	7.75	0.367	4.96
13	Adirondack Mountains	25.162	42	5.74	0.397	3.78
14	Maine West	25.857	13	14.55	0.423	1.88
15	Appalachian South	29.481	9	17.43	0.351	3.38
16	Minnesota North Woods	39.831	54	2.66	0.42	1.69
17	Wisconsin-UP	59.462	43	2.95	0.38	2.55

Each patch identified meets the minimum thresholds for areas that could support a sufficiently large puma population to avoid genetic drift and inbreeding depression

Act in Minnesota. When prioritizing potential puma recovery areas, previously established legal protections could be regarded as an invisible component of habitat suitability.

## Discussion

Pumas are currently dispersing into the Midwest (LaRue et al. 2012), and eventually we expect there to be breeding populations in the East, though it may take considerable time. For example, it took nearly 20 years for pumas to expand beyond the Black Hills in South Dakota across 100 miles of human-dominated landscapes to establish a new breeding population in the Pine Ridge area of Nebraska (LaRue and Nielsen 2016). Patches closer to source populations in the West could be recolonized sooner than those farther away, and could serve as important sources for pumas to continue eastward recolonization.

**Table 3** A review of puma status under state endangered species law and wildlife action plans in states that overlap viable habitat patches

State	Primary wildlife authorities: agency & board	Legal status, current governing laws, & policy overview	Patch number
Arkansas	Agency: Arkansas Game and Fish Commission Board: Arkansas Game and Fish Commission	State-level endangered, threatened, or special concern species designations do not exist, and thus pumas cannot be listed as such. Not identified as a Species of Greatest Conservation Need (SGCN) in the 2015 Wildlife Action Plan (Fowler 2015)	1
Georgia	Agency: Wildlife Resources Division, Georgia Department of Natural Resources Board: Georgia Board of Natural Resources	Florida panther ( <i>Puma concolor coryi</i> ) listed as endangered under state endangered species law ( <a href="https://law.justia.com/codes/georgia/ia/2020/">https://law.justia.com/codes/georgia/ia/2020/</a> ). Also a High Priority—GA's preferred terminology for a SGCN—species in the 2015 Wildlife Action Plan (Georgia Department of Natural Resources, Wildlife Resources Division 2015). State law identifies pumas ( <i>Puma concolor</i> ) as a game species ( <a href="https://regulations.justia.com/states/georgia/">https://regulations.justia.com/states/georgia/</a> ) but prohibits any hunting season ( <a href="https://regulations.justia.com/states/georgia/">https://regulations.justia.com/states/georgia/</a> )	10
Kentucky	Agency: Kentucky Department of Fish & Wildlife Resources, Kentucky Tourism, Arts and Heritage Cabinet Board: Kentucky Fish and Wildlife Commission	Not listed under state endangered species law (Kentucky Department of Fish & Wildlife Resources 2022), not identified as a SGCN in the 2015 Wildlife Action Plan (Kentucky Department of Fish and Wildlife Resources 2013)	15
Maine	Agency: Maine Department of Inland Fisheries & Wildlife	Not listed under state endangered species law (Maine Department of Inland Fisheries and Wildlife 2015a, b), not identified as a SGCN in the 2015 Wildlife Action Plan (Maine Department of Inland Fisheries and Wildlife 2015a, b)	8, 14
Massachusetts	Agency: Division of Fisheries & Wildlife, Massachusetts Department of Fish and Game Board: Massachusetts Fisheries and Wildlife Board	Not listed under state endangered species law ( <a href="https://regulations.justia.com/states/massachusetts/">https://regulations.justia.com/states/massachusetts/</a> ), not identified as a SGCN in the 2015 Wildlife Action Plan (Massachusetts Division of Fisheries and Wildlife 2015)	2, 5
Michigan	Agency: Michigan Department of Natural Resources Board: Michigan Natural Resources Commission	Puma ( <i>Puma concolor</i> ) listed as endangered under state endangered species law ( <a href="https://regulations.justia.com/states/michigan/">https://regulations.justia.com/states/michigan/</a> ), not identified as a SGCN in the 2015 Wildlife Action Plan (Michigan Department of Natural Resources 2015)	3, 4

**Table 3** (continued)

State	Primary wildlife authorities: agency & board	Legal status, current governing laws, & policy overview	Patch number
Minnesota	Agency: Minnesota Fish and Wildlife Division, Department of Natural Resources	Puma ( <i>Puma concolor</i> ) listed as a species of special concern under state endangered species law ( <a href="https://regulations.justia.com/states/minnesota/">https://regulations.justia.com/states/minnesota/</a> ), but that status does not confer the protection of endangered or threatened species( <a href="https://law.justia.com/codes/minnesota/2021/">https://law.justia.com/codes/minnesota/2021/</a> ). However, it does automatically make it a SGCN in the 2015 Wildlife Action Plan(Minnesota Department of Natural Resources 2015) No open season for pumas( <a href="https://law.justia.com/codes/minnesota/2021/">https://law.justia.com/codes/minnesota/2021/</a> ), and not on the list of animals that may be killed if causing damage( <a href="https://law.justia.com/codes/minnesota/2021/">https://law.justia.com/codes/minnesota/2021/</a> )	16
Missouri	Agency: Missouri Department of Conservation Board: Missouri Conservation Commission	Not listed under state endangered species law(Missouri Department of Conservation 2021), not identified as a SGCN in the 2015 Wildlife Action Plan(Missouri Department of Conservation (2015)	9
New Hampshire	Agency: New Hampshire Fish and Game Department Board: New Hampshire Fish and Game Commission	Not listed under state endangered species law( <a href="https://regulations.justia.com/states/new-hampshire/">https://regulations.justia.com/states/new-hampshire/</a> ), not identified as a SGCN in the 2015 Wildlife Action Plan( <a href="https://www.wildlife.state.nh.us/wildlife/wap.html">https://www.wildlife.state.nh.us/wildlife/wap.html</a> ). Taking or killing mountain lions is prohibited, except in cases of self or property protection( <a href="https://law.justia.com/codes/new-hampshire/2021/">https://law.justia.com/codes/new-hampshire/2021/</a> ); however, the director of Fish & Game may take measures they deem necessary for controlling this species should they “become a nuisance in any part of the state.” ( <a href="https://law.justia.com/codes/new-hampshire/2021/">https://law.justia.com/codes/new-hampshire/2021/</a> )	2
New York	Agency: New York Department of Environmental Conservation	Puma ( <i>Puma concolor</i> ) listed as endangered under state endangered species law( <a href="https://regulations.justia.com/states/new-york/New-York-Current-and-Proposed-Status-of-All-Species-on-Proposed-List-;but-in-process-changes-first-drafted-in-2019-proposal-removal(https://www.dec.ny.gov/docs/wildlife_pdf/masterlistproperg.pdf">https://regulations.justia.com/states/new-york/New-York-Current-and-Proposed-Status-of-All-Species-on-Proposed-List-;but-in-process-changes-first-drafted-in-2019-proposal-removal(https://www.dec.ny.gov/docs/wildlife_pdf/masterlistproperg.pdf</a> ) Not identified as a SGCN in the 2015 Wildlife Action Plan( <a href="https://www.dec.ny.gov/docs/wildlife_pdf/swapfinaldraft2015.pdf">https://www.dec.ny.gov/docs/wildlife_pdf/swapfinaldraft2015.pdf</a> )	5, 7, 12, 13

Table 3 (continued)

State	Primary wildlife authorities: agency & board	Legal status, current governing laws, & policy overview	Patch number
North Carolina	Agency: North Carolina Wildlife Resources Commission Board: North Carolina Wildlife Resources Commission	Although the Eastern Mountain Lion ( <i>Puma concolor cougar</i> ) was removed from the federal endangered species list in 2018, as of an October 1st, 2021 update to North Carolina's endangered species list, <i>Puma concolor</i> was still identified as federally endangered in North Carolina (North Carolina Administrative Code (2021)) Not identified as a SGCN in the 2015 Wildlife Action Plan (North Carolina Wildlife Resources Commission (2015))	10
Pennsylvania	Agency: Pennsylvania Game Commission Board: Pennsylvania Board of Game Commissioners	Not listed under state endangered species law ( <a href="https://regulations.justia.com/states/pennsylvania/">https://regulations.justia.com/states/pennsylvania/</a> ), not identified as a SGCN in the 2015 Wildlife Action Plan (Pennsylvania Game Commission (2015))	7, 12
South Carolina	Agency: Wildlife and Freshwater Fisheries Division, South Carolina Department of Natural Resources Board: South Carolina Department of Natural Resources Board	Eastern Mountain Lion ( <i>Puma concolor cougar</i> ) listed as endangered under state endangered species law (South Carolina Code of Regulations (2022)). Not identified as a SGCN in the 2015 Wildlife Action Plan (South Carolina Department of Natural Resources (2015))	10
Tennessee	Agency: Tennessee Wildlife Resources Agency Board: Tennessee Fish & Wildlife Commission	Not listed under state endangered species law (Tennessee Comp Rules and Regs (2022)), not identified as a SGCN in the 2015 Wildlife Action Plan (Tennessee Wildlife Resources Agency (2015))	10, 15
Vermont	Agency: Department of Fish & Wildlife, Agency of Natural Resources Board: Vermont Fish & Wildlife Board	Eastern Mountain Lion ( <i>Puma concolor cougar</i> ) listed as endangered under state endangered species law ( <a href="https://regulations.justia.com/states/vermont/">https://regulations.justia.com/states/vermont/</a> ), and is identified as a SGCN of medium (as opposed to high) priority in the 2015 Wildlife Action Plan (Vermont Fish and Wildlife Department 2015)	5
West Virginia	Agency: Wildlife Resources Section, Division of Natural Resources, West Virginia Department of Commerce	State-level endangered, threatened, or special concern species designations do not exist, and thus pumas cannot be listed as such. Not identified as a SGCN in the 2015 Wildlife Action Plan (West Virginia Division of Natural Resources 2015)	6, 11, 15

**Table 3** (continued)

State	Primary wildlife authorities: agency & board	Legal status, current governing laws, & policy overview	Patch number
Wisconsin	Agency: Fish, Wildlife & Parks Division, Wisconsin Department of Natural Resources Board: Wisconsin Natural Resources Board	Not listed under state endangered species law( <a href="https://regulations.justia.com/states/wisconsin/">https://regulations.justia.com/states/wisconsin/</a> ), not identified as a SGCN in the 2015 Wildlife Action Plan(Wisconsin Department of Natural Resources (2015). Pumas are a protected wild animal, which means essentially any taking, attempted taking, transport, or possession of them alive or dead must be explicitly authorized by the Department of Natural Resources( <a href="https://regulations.justia.com/states/wisconsin/">https://regulations.justia.com/states/wisconsin/</a> ))	17

Additional puma-related laws and regulations were also identified; this list is not exhaustive but includes those most directly related to puma protection in these states. Note, the state wildlife board is only included if the board has any degree of regulatory authority

For habitat to be considered suitable for pumas, there needs to be appropriate vegetation communities with low human use, livestock use, and highway density. Our analyses highlighted numerous large habitat patches that could support sustainable puma populations in the East. Of course, as the human footprint continues to expand, the suitable habitat we identified may shrink, reducing the number of places where puma populations can maintain genetic health without conservation interventions. For example, our analyses did not identify southern Florida as suitable habitat to accommodate sustainable puma populations, even though it currently hosts the only breeding population of pumas in the East. However, a recent analysis has revealed that this population will require further conservation support via the translocation of pumas from elsewhere to boost the genetic diversity among Florida pumas (locally referred to as panthers) to maintain a viable puma population into the future (van de Kerk et al. 2019).

Interstates played a large role in separating potential puma habitat patches in our results, especially in the New England states (Fig. 2a and b): Vermont, New Hampshire, Massachusetts, and Maine. There are a few instances in which adjacent areas would be considered one large patch if they were not separated by interstates (e.g., Michigan West and Michigan East; Allegheny Plateau, Appalachian North, and Appalachian South; Fig. 2). Dispersing pumas, which carry vital genetic material between habitat patches and puma populations and help restore genetic health, will attempt to cross major highways and may even be successful sometimes (e.g., Hawley et al. 2016). Therefore, many of the divisions between patches in our results may in fact be somewhat permeable from a gene flow perspective, and what appears as several smaller patches could be connected to create fewer, larger patches. Our analyses treated interstates as uniform across their length, but a better metric to determine what sections of specific interstates are and are not permeable to pumas would be traffic volume. For example, Interstate 93, which runs north–south in New Hampshire, has very low traffic volume in the north and higher use in the south, so we would expect the southern portion to be less permeable to pumas than the northern portion. Unfortunately, we currently lack specific threshold information about what traffic loads restrict pumas and impact their metapopulation dynamics (e.g., Gustafson et al. 2019).

We did not rank patches in terms of potential suitability for pumas, because they all met the conservation thresholds we set and therefore all could host puma populations in the future. Nevertheless, the metrics we used to delineate these areas, and those we summarize in Table 2, may be useful for conservation planning comparison. For example, the larger the size of an area of suitable habitat, generally the greater chance that a resident flora or fauna population will remain viable (Bender et al. 1998). The lowest human and livestock densities might be useful in identifying areas where pumas are least likely to experience conflict with human communities, and the highest sociocultural index values (Manfredo et al. 2021) may be indicative of potential tolerance and the willingness of local people to coexist with large carnivores.

Pumas play an important role in regulating prey, producing carrion for scavengers, and providing ecosystem services, each of which contributes to ecological resilience and healthy human communities (Barry et al. 2019; LaBarge et al. 2022). Pumas may more directly benefit humans by reducing automobile collisions with deer, protecting human lives and reducing expensive damages (Gilbert et al. 2021). However, these and other environmental benefits are unlikely to convince all members of the public to support puma recovery. Education and outreach will play an essential role in communicating the potential costs and benefits of living sympatrically with large carnivores as they recolonize their historic range (López-Bao et al. 2017; Gilbert et al. 2021).



Pumas pose real and perceived risks to domestic animals and humans, and so our results also serve as a map to identify priority areas for investing in outreach, education, and policy instruments before pumas reestablish in the East (Madden 2004; Baruch-Mordo et al. 2011; Redpath et al. 2013). While some details of human-puma conflict may be unpredictable in areas newly colonized by pumas, important patterns of conflict are readily inferred from those documented following other carnivore restoration or recolonization, such as gray wolves (*Canis lupus*) in North America and Europe. For example, there is lively research debate and public discord surrounding the relationship between wolf recovery and hunter opportunity (Garrot et al. 2005; Vucetich et al. 2005; Brodie et al. 2013; MacNulty et al. 2016; Wikenros et al. 2020). Livestock-carnivore conflict and how to manage reestablished carnivore populations on rangelands are similarly contentious topics (e.g., lethally removing wolves to protect livestock; Bradley et al. 2015; Santiago-Avila et al. 2018).

Anticipating conflict and finding ways to get ahead of issues common to other carnivore recovery efforts could help facilitate puma recovery. Framing this in the context of wolf restoration, public acceptance of wolves decreased among some stakeholders following their return (Ericsson and Heberlein 2003; Treves et al. 2013), even as broader public attitudes (e.g., US, Sweden) became more positive (George et al. 2016; Heberlein and Ericsson 2008). Although some have suggested that decreased acceptance of wolves within an area tends to accompany their occurrence (e.g., Karlsson and Sjöström, 2007), recent analyses in the U.S. show this effect depends upon the stakeholder groups with which one identifies (Carlson et al. 2020). Specifically, a large-scale survey of Americans residing within and outside wolf recovery areas found ranchers who lived within those areas were more negative towards wolves than ranchers who lived outside of those areas; conversely, environmentalists living within wolf recovery areas were actually more positive towards wolves than environmentalists living outside these areas. While it may be tempting to assume similar social conflict would follow puma restoration, differences in the behavior and ecology of pumas and wolves (e.g., ambush vs. coursing predation; cryptic vs. visible behavior, etc.) may lead to pumas being perceived as less threatening to livestock and game populations. Likewise, regional differences in values (see Manfredo et al. 2020) and political ideologies change the cultural context in which conflicts with carnivores are perceived. Thus, we caution readers against assuming the type of conflict that has accompanied wolf restoration efforts will also accompany puma restoration.

Our analyses suggest that ample habitat exists in the eastern US to support pumas, and ongoing records of puma dispersal in the Midwest show that individuals are on their way. Future analyses could help predict how long it might take to see breeding pumas establish in the east. The question remains, however, whether the people of the East are willing to coexist with pumas when they arrive. We suggest proactive efforts (as opposed to reactive, post-problem management) to help residents avoid negative impacts of pumas, will be the most useful for promoting coexistence.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s10531-022-02529-z>.

**Acknowledgements** We thank Tompkins Conservation, the Donald Slavik Foundation, the Kisco Cares Foundation, the Ayers Wildcat Conservation Trust, D. McFarland and the Bromley Charitable Trust for supporting and funding this work.

**Author contributions** All authors contributed to the study conception, design, data collection, analyses, manuscript writing, and/or revisions. All authors read and approved the final manuscript.

**Funding** This work was supported by Tompkins Conservation, the Donald Slavik Foundation, the Kisco Cares Foundation, Bromley Charitable Trust, and the Ayers Wildcat Conservation Trust. The authors have no relevant financial or non-financial interests to disclose. Ayers Wildcat Conservation Trust, Tompkins Conservation, Donald Slavik Foundation, Bromley Charitable Trust, Kisco Cares Foundation

**Data availability** Each of the datasets analyzed in this study are available from the sources described in Table 1.

## Declarations

**Competing interest** The authors have no competing interests as defined by Springer, or other interests that might be perceived to influence the results and/or discussion reported in this paper.

**Ethical approval** No institutional approval or ethical review was needed in order to conduct this study.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Barry JM, Elbroch LM, Aiello-Lammens ME, Sarno RJ, Seelye L, Kusler A, Quigley HB, Grigione MM (2019) Pumas as ecosystem engineers: ungulate carcasses support beetle assemblages in the Greater Yellowstone Ecosystem. *Oecologia* 189(3):577–586. <https://doi.org/10.1007/s00442-018-4315-z>
- Baruch-Mordo S, Breck SW, Wilson KR, Broderick J (2011) The carrot or the stick? evaluation of education and enforcement as management tools for human-wildlife conflicts. *PLoS ONE* 6(1):e15681
- Baune C, Wolfe LL, Schott KC, Griffin KA, Hughson AG, Miller MW, Race B (2021) Reduction of Chronic Wasting Disease prion seeding activity following digestion by mountain lions. *Mosphere* 6(6):e00812-e821
- Bender DJ, Contreras TA, Fahrig L (1998) Habitat Loss and Population Decline: A Meta-Analysis of the Patch Size Effect. *Ecology* 79:517–533
- Bradley EH, Robinson HS, Bangs EE, Kunkel K, Jimenez MD, Gude JA, Grimm T (2015) Effects of Wolf Removal on Livestock Depredation Recurrence and Wolf Recovery in Montana, Idaho, and Wyoming. *J Wildl Manag* 79:1337–1346
- Brocke, R.H., (1981). Reintroduction of the Cougar *Felis Concolor* in Adirondack Park: A Problem Analysis and Recommendations. Adirondack Wildlife Research Project Reports Funded by the Pittman-Robertson Act. 9. <https://digitalcommons.esf.edu/awrpr/9>
- Brodie JF, Johnson H, Mitchell M, Zager P, Proffitt K, Hebblewhite M, Kauffman M, Johnson B, Bissonette J, Bishop C, Gude J, Herbert J, Hersey K, Hurley M, Lukacs p, McCorquodale S, McIntire E, Nowak J, Sawyer H, Smith D, White PJ (2013) Relative influence of human harvest, carnivores, and weather on adult female elk survival across western North America. *J Appl Ecol* 50:295–305
- Burdett CL, Crooks KR, Theobald DM, Wilson KR, Boydston EE, Lyren LM, Fisher RN, Vickers TW, Morrison SA, Boyce WM (2010) Interfacing models of wildlife habitat and human development to predict the future distribution of puma habitat. *Ecosphere* 1(1):1–21. <https://doi.org/10.1890/ES10-00005.1>
- California Department of Fish and Wildlife (2015) Report to the fish and game commission regarding findings of necropsies on mountain lions taken under depredation permits in 2015. State of California Natural Resources Agency Department of Fish and Wildlife. p 3
- Carlson SC, Dietsch AM, Slagle KM, Bruskotter JT (2020) The VIPs of wolf conservation: How values, identity, and place shape attitudes toward wolves in the United States. *Front Ecol Evol* 8:6

- Center for International Earth Science Information Network - CIESIN - Columbia University. (2018). Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 11. Palisades, New York: NASA Socioeconomic Data and Applications Center (SEDAC). doi.org/<https://doi.org/10.7927/H49C6VHW>. Accessed January 15, 2022.
- Compilation of Rules and Regulations of the State of Georgia. GA Code Ann., § 27–1–2. Last modified April 4, 2022. <https://regulations.justia.com/states/georgia/>. Accessed 8 June 2022
- Compilation of Rules and Regulations of the State of Georgia. GA Code Ann., § 27–3–15. Last modified April 4, 2022. <https://regulations.justia.com/states/georgia/>. Accessed 8 June 2022
- Culver M, Johnson WE, Pecon-Slattery J, O'Brien SJ (2000) Genomic ancestry of the American Puma (*Puma concolor*). *J Hered* 91:186–197
- Danz HP (1999) *Cougar!* Swallow Press, Athens, Ohio, p 310
- Dellinger JA, Gustafson KD, Gammons DJ, Ernest HB, Torres SG (2020) Minimum habitat thresholds required for conserving mountain lion genetic diversity. *Ecol Evol* 10(19):10687–10696. <https://doi.org/10.1002/ece3.6723>
- Dickson BG, Jenness JS, Beier P (2005) Influence of Vegetation, Topography, and Roads on Cougar Movement in Southern California. *J Wildl Manag* 69(1):264–276. [https://doi.org/10.2193/0022-541x\(2005\)069%3c0264:iovtar%3e2.0.co;2](https://doi.org/10.2193/0022-541x(2005)069%3c0264:iovtar%3e2.0.co;2)
- EPA 2022. Ecoregions of North America. <https://www.epa.gov/eco-research/ecoregions-north-america> Accessed Spetember 8, 2022.
- Ericsson G, Heberlein TA (2003) Attitudes of hunters, locals, and the general public in Sweden now that the wolves are back. *Biol Cons* 111:149–159
- Feldman JW (2007) Public opinion, the Leopold Report, and the reform of federal predator control policy. *Human-Wildlife Conflicts* 1(1):112–124
- Fowler, A, ed. (2015) *Arkansas Wildlife Action Plan*. Little Rock, AR. <https://www.wildlifearkansas.com/strategy.html>. Accessed 8 June 2022
- Gantchoff MG, Erb JD, MacFarland DM, Norton DC, Price Tack JL, Roell BJ, Belant JL (2021) Potential distribution and connectivity for recolonizing cougars in the Great Lakes region, USA. *Biol Cons* 257:109144
- Garrott, R.A., Gude, J.A., Bergman, E.J., Gower, C., White, P.J. and Hamlin, K.L. (2005) Generalizing wolf effects across the Greater Yellowstone Area: a cautionary note. *Wildl Soc Bull* 33(4):1245–1255
- Georgia Department of Natural Resources, Wildlife Resources Division (2015) *Georgia State Wildlife Action Plan*. Social Circle, GA. [https://georgiawildlife.com/sites/default/files/wrd/pdf/swap/SWAP2015MainReport\\_92015.pdf](https://georgiawildlife.com/sites/default/files/wrd/pdf/swap/SWAP2015MainReport_92015.pdf). Accessed 8 June 2022
- George KA, Slagle KM, Wilson RS, Moeller SJ, Bruskotter JT (2016) Changes in attitudes toward animals in the United States from 1978 to 2014. *Biol Cons* 201:237–242
- Georgia Rules and Regulations 391–4–10-.09. Georgia Code. Last modified 2020. <https://law.justia.com/codes/georgia/2020/>. Accessed 8 June 2022
- Gilbert S, Carter N, Naidoo R (2021) Predation services: Quantifying societal effects of predators and their prey. *Front Ecol Environ* 19(5):292–299
- Gill, R.B. (2010) “To save a mountain lion: evolving philosophy of nature and cougars,” in *Cougar Ecology and Conservation* (Ed Mornocker M and Negri S). pp 5–16. University of Chicago Press
- Glick HB (2014) Modeling cougar habitat in the Northeastern United States. *Ecol Model* 285:78–89
- Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R (2017) Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sens Environ* 202:18–27
- Gray M, Wilmers CC, Reed SE, Merenlender AM (2016) Landscape feature-based permeability models relate to puma occurrence. *Landsc Urban Plan* 147:50–58
- Guerisoli MDLM, Luengos Vidal E, Caruso N, Giordano AJ, Lucherini M (2021) Puma–livestock conflicts in the Americas: A review of the evidence. *Mammal Rev* 51(2):228–246
- Gustafson KD, Gagne RB, Vickers TW, Riley SP, Wilmers CC, Bleich VC, Pierce BM, Kenyon M, Drzenovich TL, Sikich JA, Boyce WM (2019) Genetic source–sink dynamics among naturally structured and anthropogenically fragmented puma populations. *Conserv Genet* 20(2):215–227
- Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, Tyukavina A, Thau D, Stehman SV, Goetz SJ, Loveland TR, Kommareddy A (2013) High-resolution global maps of 21st-century forest cover change. *Science* 342(6160):850–853
- Hawley JE, Rego PW, Wydeven AP, Schwartz MK, Viner TC, Kays R, Pilgrim KL, Jenks JA (2016) Long-distance dispersal of a subadult male cougar from South Dakota to Connecticut documented with DNA evidence. *J Mammal* 97(5):1435–1440
- Heberlein TA, Ericsson G (2008) Public attitudes and the future of wolves *Canis lupus* in Sweden. *Wildl Biol* 14(3):391–394

- Karlsson J, Sjöström M (2007) Human attitudes towards wolves, a matter of distance. *Biol Cons* 137(4):610–616
- Kentucky Department of Fish and Wildlife Resources (2013) Kentucky's Comprehensive Wildlife Conservation Strategy. Frankfort, KY. <http://fw.ky.gov/WAP/Pages/Default.aspx>. Accessed 8 June 2022
- Kentucky Department of Fish & Wildlife Resources (2022) Species Information. <https://app.fw.ky.gov/speciesinfo/speciesList.asp?strGroup=11&strSort1=Class&strSort2=CommonName>. Accessed 8 June 2022
- Knopff AA, Knopff KH, Boyce MS, St. Clair, C.C. (2014) Flexible habitat selection by cougars in response to anthropogenic development. *Biol Cons* 178:136–145. <https://doi.org/10.1016/j.biocon.2014.07.017>
- LaBarge LR, Evans MJ, Miller JRB, Cannataro G, Hunt C, Elbroch LM (2022) Pumas *Puma concolor* as ecological brokers: a review of their biotic relationships. *Mammal Rev*. <https://doi.org/10.1111/mam.12281>
- LaRue MA, Nielsen CK (2008) Modelling potential dispersal corridors for cougars in midwestern North America using least-cost path methods. *Ecol Model* 212(3–4):372–381
- LaRue MA, Nielsen CK (2011) Modelling potential habitat for cougars in midwestern North America. *Ecol Model* 222:897–900
- LaRue MA, Nielsen CK (2016) Population viability of recolonizing cougars in midwestern North America. *Ecol Modell* 321(2016):121–129
- LaRue MA, Nielsen CK, Dowling M, Miller K, Wilson B, Shaw H, Anderson CR Jr (2012) Cougars are recolonizing the midwest: analysis of cougar confirmations during 1990–2008. *J Wildl Manag* 76(7):1364–1369
- LaRue MA, Nielsen CK, Pease BS (2022) Increases in Midwestern cougars despite harvest in a source population. *Jour Wild Mgmt* 83:1306–1313
- Laundré JW (2013) The feasibility of the north-eastern USA supporting the return of the cougar *Puma concolor*. *Oryx* 47(1):96–104
- Laundré JW, Hernández L (2003) Winter hunting habitat of pumas *Puma concolor* in northwestern Utah and southern Idaho, USA. *Wildl Biol* 9(4):123–129
- López-Bao JV, Bruskotter J, Chapron G (2017) Finding space for large carnivores. *Nature, Ecology, and Evolution* 1(5):1–2
- MacNulty DR, Stahler DR, Wyman CT, Ruprecht J, Smith DW (2016) The challenge of understanding northern Yellowstone elk dynamics after wolf reintroduction. *Yellowstone Science* 24(1):25–33
- Madden F (2004) Creating coexistence between humans and wildlife: global perspectives on local efforts to address human–wildlife conflict. *Hum Dimens Wildl* 9(4):247–257
- Maine Department of Inland Fisheries and Wildlife (2015a) State List of Endangered & Threatened Species. Last modified October 15, 2015a. <https://www.maine.gov/ifw/fish-wildlife/wildlife/endangered-threatened-species/listed-species.html>. Accessed 8 June 2022
- Maine Department of Inland Fisheries and Wildlife (2015b) *Maine's Wildlife Action Plan*. Augusta ME. [https://www.maine.gov/ifw/docs/swap/2015b\\_MESWAP.pdf](https://www.maine.gov/ifw/docs/swap/2015b_MESWAP.pdf). Accessed 8 June 2022
- Manfredo MJ, Berl REW, Teel TL, Bruskotter JT (2021) Bringing social values to wildlife conservation decisions. *Front Ecol Environ* 19(6):355–362
- Massachusetts Division of Fisheries and Wildlife (2015). Massachusetts State Wildlife Action Plan. Westborough, MA. <https://www.mass.gov/files/documents/2016/12/sy/ma-swap-public-draft-26june2015-chapter1.pdf>. Accessed 8 June 2022
- Massachusetts Regulations 321 CMR 10.90. Last modified April 1, 2022. <https://regulations.justia.com/states/massachusetts/>. Accessed 8 June 2022
- Mattson, D. and S. Clark (2010). "People, politics and cougar management," in *Cougar Ecology and Conservation* (Ed Mornocker M and Negri S). pp 206–220. University of Chicago Press
- McCollough, M. (2011). Eastern puma (cougar) (*Puma concolor cougar*) 5-year review: summary and evaluation. 110p.
- Michigan Department of Natural Resources (2015) Michigan's Wildlife Action Plan. Lansing, MI. [https://www.michigan.gov/dnr/-/media/Project/Websites/dnr/Documents/WLD/WAP/18\\_appendix2\\_sgcn\\_rationales.pdf?rev=186c9c133a744cf1848f3260395bf6e4](https://www.michigan.gov/dnr/-/media/Project/Websites/dnr/Documents/WLD/WAP/18_appendix2_sgcn_rationales.pdf?rev=186c9c133a744cf1848f3260395bf6e4). Accessed 8 June 2022
- Michigan Administrative Code R. 299.1027 Last modified April 1, 2022. <https://regulations.justia.com/states/michigan/>. Accessed 8 June 2022
- Minnesota Administrative Rules. 6134.0150. Last modified 2021. <https://law.justia.com/codes/minnesota/2021/>. Accessed 8 June 2022
- Minnesota Administrative Rules. 6134.0200. Last modified 2022. <https://regulations.justia.com/states/minnesota/>. Accessed 8 June 2022





- Minnesota Department of Natural Resources (2015) Minnesota's Wildlife Action Plan, 2015–2025. St. Paul, MN. <https://files.dnr.state.mn.us/assistance/nrplanning/bigpicture/mnwap/wildlife-action-plan-2015-2025.pdf>. Accessed 8 June 2022
- Minnesota Statutes. § 97B.641. Last modified 2021a. <https://law.justia.com/codes/minnesota/2021a/>. Accessed 8 June 2022
- Minnesota Statutes. § 97B.655. Last modified 2021b. <https://law.justia.com/codes/minnesota/2021b/>. Accessed 8 June 2022
- Missouri Department of Conservation (2015) Missouri State Wildlife Action Plan. Jefferson City, MO. [https://mdc.mo.gov/sites/default/files/2020-04/SWAP\\_0.pdf](https://mdc.mo.gov/sites/default/files/2020-04/SWAP_0.pdf). Accessed 8 June 2022
- Missouri Department of Conservation (2021). Missouri Species and Communities of Conservation Concern Checklist. Jefferson City, MO: Missouri Department of Conservation. [https://education.mdc.mo.gov/sites/default/files/downloads/2021\\_SOCC.pdf](https://education.mdc.mo.gov/sites/default/files/downloads/2021_SOCC.pdf). Accessed 8 June 2022
- New Hampshire Code of Administrative Rules. NH Admin Rules Fis 1001.01, NH Admin Rules Fis 1001.02. Last modified April 7, 2022. <https://regulations.justia.com/states/new-hampshire/>. Accessed 8 June 2022
- New Hampshire Department of Fish and Game (2015) New Hampshire Wildlife Action Plan. Concord, NH. <https://www.wildlife.state.nh.us/wildlife/wap.html>. Accessed 8 June 2022
- New Hampshire Revised Statutes. NH Rev. Stat. § 208:1-b Last modified 2021c. <https://law.justia.com/codes/new-hampshire/2021c/>. Accessed 8 June 2022
- New Hampshire Revised Statutes. NH Rev. Stat. § 208:1-c Last modified 2021. <https://law.justia.com/codes/new-hampshire/2021/>. Accessed 8 June 2022
- New York Codes, Rules and Regulations. 6 NY Comp Codes Rules and Regs § 182.5. Last modified April 6, 2022. <https://regulations.justia.com/states/new-york/New York Current and Proposed Status of All Species on Proposed List>. Accessed June 12, 2022.
- New York Department of Environmental Conservation (2015) New York State Wildlife Action Plan. Albany, NY. [https://www.dec.ny.gov/docs/wildlife\\_pdf/swapfinaldraft2015.pdf](https://www.dec.ny.gov/docs/wildlife_pdf/swapfinaldraft2015.pdf). Accessed 8 June 2022
- Nielsen C, Thompson D, Kelly M, Lopez-Gonzalez CA (2017) *Puma concolor*, puma. IUCN Red List of Threatened Species 8235:12
- North Carolina Administrative Code (2021) 15A NC Admin. Code 10I.0103. Protected Wildlife Species of North Carolina. Last modified October 1, 2021. <https://www.ncwildlife.org/Portals/0/Conserving/documents/Protected-Wildlife-Species-of-NC.pdf>. Accessed 8 June 2022
- North Carolina Wildlife Resources Commission (2015) North Carolina Wildlife Action Plan. Raleigh, NC. <https://www.ncwildlife.org/Portals/0/Conserving/documents/2015WildlifeActionPlan/NC-WAP-2015-All-Documents.pdf>. Accessed 8 June 2022
- O'Neil S.T., Rahn K.C., Bump J.K. (2014) Habitat Capacity for Cougar Recolonization in the Upper Great Lakes Region. *PLoS ONE* 9(11):e112565
- Pennsylvania Game Commission (2015) Pennsylvania Wildlife Action Plan. Harrisburg, PA. <https://www.pgc.pa.gov/Wildlife/WildlifeActionPlan/Pages/default.aspx>. Accessed 8 June 2022
- Pennsylvania Code. 58 PA Code § 133.41. Last modified April 16, 2022. <https://regulations.justia.com/states/pennsylvania/>. Accessed 8 June 2022
- Poole KG, Mowat G (2005) Winter habitat relationships of deer and elk in the temperate interior mountains of British Columbia. *Wildl Soc Bull* 33(4):1288–1302
- Pritchard, J.A., (2021). The American Society of Mammalogists, The Ecological Society of America, and the Politics of Preservation. *Studies in the History of Biology*, 13(2).
- Redpath SM, Young J, Evely A, Adams WM, Sutherland WJ, Whitehouse A, Amar A, Lambert RA, Linnell JD, Watt A, Gutierrez RJ (2013) Understanding and managing conservation conflicts. *Trends Ecol Evol* 28(2):100–109
- Ripple, W. J., Estes, J. A., Beschta, R. L., Wilmers, C. C., Ritchie, E. G., Hebblewhite, M., Berger, J., Elmhagen, B., Letnic, M., Nelson, M. P., Schmitz, O. J., Smith, D. W., Wallach, A. D., Wirsing, A. J. (2014). Status and ecological effects of the world's largest carnivores. *Science*, 343(6167). doi.org/<https://doi.org/10.1126/science.1241484>
- Robinson TP, Wint GW, Conchedda G, Van Boeckel TP, Ercoli V, Palamara E, Cinardi G, D'Aiotti L, Hay SI, Gilbert M (2014) Mapping the global distribution of livestock. *PLoS One* 9(5):e96084
- Santiago-Avila FJ, Cornman AM, Treves A (2018) Killing wolves to prevent predation on livestock may protect one farm but harm neighbors. *PLoS ONE* 13:e0189729
- Smith JB, Nielsen CK, Helligren EC (2016) Suitable habitat for recolonizing large carnivores in the mid-western USA. *Oryx* 50:555–564

- South Carolina Code of Regulations (2022) SC Code Regs 123–150. Last modified March 25, 2022. <https://regulations.justia.com/states/south-carolina/chapter-123/article-5/section-123-150/>. Accessed 8 June 2022
- South Carolina Department of Natural Resources (2015) South Carolina's State Wildlife Action Plan. Columbia, SC. <https://www.dnr.sc.gov/swap/main/2015StateWildlifeActionPlan-chaptersonly.pdf>. Accessed 8 June 2022
- Sulla-Menashe D, Friedl MA (2018) User guide to collection 6 MODIS land cover (MCD12Q1 and MCD12C1) product. USGS, Reston, VA, USA, 1, p 18
- Sweaner LL, Logan KA, Hornocker MG (2000) Cougar dispersal patterns, metapopulation dynamics, and conservation. *Conserv Biol* 14:798–808
- Tennessee Comp Rules and Regs (2022) 1660–01–32-.02. Last modified May 9, 2022. <https://regulations.justia.com/states/tennessee/>. Accessed 8 June 2022
- Tennessee Wildlife Resources Agency (2015) Tennessee Wildlife Action Plan. Nashville, TN. <http://twraonline.org/2015swap.pdf>. Accessed 8 June 2022
- The New York State Department of Environmental Conservation (2020) Current and Proposed Status of All Species on Proposed List. [https://www.dec.ny.gov/docs/wildlife\\_pdf/masterlistpropreg.pdf](https://www.dec.ny.gov/docs/wildlife_pdf/masterlistpropreg.pdf). Accessed June 12, 2022
- Treves A, Naughton-Treves L, Shelley V (2013) Longitudinal analysis of attitudes toward wolves. *Conserv Biol* 27(2):315–323
- van de Kerk M, Onorato DP, Hostetler JA, Bolker BM, Oli MK (2019) Dynamics, persistence, and genetic management of the endangered Florida Panther population. *Wildl Monogr* 203:3–35
- Vermont Code of Rules. 12 010 012. Last modified March 2022. <https://regulations.justia.com/states/vermont/>. Accessed 8 June 2022
- Vermont Fish and Wildlife Department (2015) Vermont's Wildlife Action Plan. Montpelier, VT: Vermont Fish and Wildlife Department. [https://vtfishandwildlife.com/sites/fishandwildlife/files/documents/About%20Us/Budget%20and%20Planning/WAP2015/\\_2015-VT-Wildlife-Action-Plan.pdf](https://vtfishandwildlife.com/sites/fishandwildlife/files/documents/About%20Us/Budget%20and%20Planning/WAP2015/_2015-VT-Wildlife-Action-Plan.pdf). Accessed 8 June 2022
- Vucetich JA, Smith DW, Stabler DR (2005) Influence of harvest, climate and wolf predation on Yellowstone elk, 1961–2004. *Oikos* 111:259–270
- West Virginia Division of Natural Resources (2015) West Virginia State Wildlife Action Plan. South Charleston, WV. <https://wvdnr.gov/wp-content/uploads/2021/05/2015-West-Virginia-State-Wildlife-Action-Plan-Submittal-1.pdf>. Accessed 8 June 2022
- Wikenros C, Sand H, Månsson J, Maartmann E, Eriksen A, Wabakken P, Zimmermann B (2020) Impact of a Recolonizing, Cross-Border Carnivore Population on Ungulate Harvest in Scandinavia. *Sci Rep* 10:21670
- Wilmers, C.C., Wang, Y., Nickel, B., Houghtaling, P., Shakeri, Y., Allen, M.L., Kermish-Wells, J., Yovovich, V., Williams, T. (2013). Scale Dependent Behavioral Responses to Human Development by a Large Predator, the Puma. *PLoS One*, 8(4). doi.org/<https://doi.org/10.1371/journal.pone.0060590>
- Wisconsin Administrative Code. § NR 27.03. Last modified March 28, 2022a. <https://regulations.justia.com/states/wisconsin/>. Accessed 8 June 2022a
- Wisconsin Administrative Code. § NR 10.02. Last modified March 28, 2022b. <https://regulations.justia.com/states/wisconsin/>. Accessed 8 June 2022b
- Wisconsin Department of Natural Resources (2015) 2015–2025 Wisconsin Wildlife Action Plan. Madison, WI. <https://p.widencdn.net/pd77jr/NH0938>. Accessed 8 June 2022
- Yang L, Sun G, Zhi L, Zhao J (2018) A new generation of the United States national land cover database: requirements, research priorities, design, and implementation strategies. *ISPRS J Photogramm Remote Sens* 146:108–123
- Yovovich V, Allen ML, Macaulay LT, Wilmers CC (2020) Using spatial characteristics of apex carnivore communication and reproductive behaviors to predict responses to future human development. *Biodivers Conserv* 29(8):2589–2603. <https://doi.org/10.1007/s10531-020-01990-y>
- Yovovich V, Thomsen M, Wilmers CC (2021) Pumas' fear of humans precipitates changes in plant architecture. *Ecosphere* 12(1). <https://doi.org/10.1002/ecs2.3309>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



## Authors and Affiliations

**Veronica Yovovich**<sup>1,2</sup>  · **Nathaniel Robinson**<sup>1,3</sup>  · **Hugh Robinson**<sup>1,4</sup>  · **Michael J. Manfredo**<sup>5</sup> · **Shelby Perry**<sup>6</sup> · **Jeremy T. Bruskotter**<sup>7</sup> · **John A. Vucetich**<sup>8</sup> · **Luis Aníbal Solórzano**<sup>9</sup> · **Lydia A. Roe**<sup>6</sup> · **Alison Lesure**<sup>10</sup> · **Jamie Robertson**<sup>1</sup> · **Tom Butler**<sup>6</sup> · **L. Mark Elbroch**<sup>1</sup> 

<sup>1</sup> Panthera, 8 West 40Th Street, 18Th Floor, New York, NY 10018, USA

<sup>2</sup> Department of Fish, Wildlife, and Conservation Biology, Colorado State University, Fort Collins, CO 80523, USA

<sup>3</sup> The Nature Conservancy, 4245 North Fairfax Drive, Suite 100, Arlington, VA 22203, USA

<sup>4</sup> Wildlife Biology Program, Department of Ecosystem and Conservation Sciences, W.A. Franke College of Forestry and Conservation, University of Montana, 32 Campus Drive, Missoula, MT 59812, USA

<sup>5</sup> Human Dimensions of Natural Resources Department, Colorado State University, Fort Collins, CO 80523, USA

<sup>6</sup> Northeast Wilderness Trust, 17 State Street, Suite 302, Montpelier, VT 05602, USA

<sup>7</sup> School of Environment and Natural Resources, Ohio State University, Columbus, Ohio 43210, USA

<sup>8</sup> College of Forest Resources and Environmental Science, Michigan Technological University, Houghton, MI 49930, USA

<sup>9</sup> Provita, Calle La Joya, Edificio Unidad Técnica del Este, Chacao, Caracas 1060, Venezuela

<sup>10</sup> Vermont Law School, 164 Chelsea St, South Royalton, Royalton, VT 05068, USA